

Received: 2015.12.09  
Accepted: 2016.01.03  
Published: 2016.01.21

e-ISSN 1643-3750  
© Med Sci Monit, 2016; 22: 226-233  
DOI: 10.12659/MSM.897044

# Differentiation of Bone Marrow Mesenchymal Stem Cells in Osteoblasts and Adipocytes and its Role in Treatment of Osteoporosis

NEW Bone

FAT of Marrow

Authors' Contribution:  
Study Design: A  
Data Collection: B  
Statistical Analysis: C  
Data Interpretation: D  
Manuscript Preparation: E  
Literature Search: F  
Funds Collection: G

ACE 1 **Cheng Wang**  
B 1 **Haoye Meng**  
DF 2 **Xin Wang**  
F 3 **Chenyang Zhao**  
A 1 **Jing Peng**  
A 1 **Yu Wang**

1 Institute of Orthopedics, Chinese PLA General Hospital, Beijing, P.R. China  
2 Department of Orthopedics, Urumqi General Hospital of Lanzhou Military Command, Urumqi, Xinjiang, P.R. China  
3 College of Life Science, University of Chinese Academy of Sciences, Beijing, P.R. China

**Corresponding Author:** Yu Wang, e-mail: wangwangdian628@163.com  
**Source of support:** This work was funded by NSFC (31240048, 31170946, 51361130034), NTRDP (2012AA020502, 2012CB518106, BWS11J025)

Osteoporosis is a systemic metabolic bone disorder characterized by a decrease in bone mass and degradation of the bone microstructure, leaving bones that are fragile and prone to fracture. Most osteoporosis treatments improve symptoms, but to date there is no quick and effective therapy. Bone marrow mesenchymal stem cells (BMMSCs) have pluripotent potential. In adults, BMMSCs differentiate mainly into osteoblasts and adipocytes in the skeleton. However, if this differentiation is unbalanced, it may lead to a decrease in bone mass. If the number of adipocyte cells increases and that of osteoblast cells decreases, osteoporosis can result. A variety of hormones and cytokines play an important role in the regulation of BMMSCs bidirectional differentiation. Therefore, a greater understanding of the regulation mechanism of BMMSC differentiation may provide new methods to prevent and treat osteoporosis. In addition, autologous, allogeneic BMMSCs or genetically modified BMMSC transplantation can effectively increase bone mass and density, increase bone mechanical strength, correct the imbalance in bone metabolism, and increase bone formation, and is expected to provide a new strategy and method for the treatment of osteoporosis.

**MeSH Keywords:** Adipocytes • Cell Transplantation • Osteoblasts • Osteoporosis • Stem Cell Research

**Full-text PDF:** <http://www.medscimonit.com/abstract/index/idArt/897044>

4159 — 83





## Background

Osteoporosis is a common metabolic bone disease characterized by decreases in bone mass and density and degradation of the bone microstructure, resulting in fragile bones that are prone to fracture. This process is the same for primary and secondary osteoporosis [1]. With an aging population, the incidence of osteoporosis is increasing each year. The latest statistics show that the number of osteoporosis patients in China has reached 90 million, and in women older than 50 years the incidence is greater than 20% [2]. Osteoporosis causes a higher incidence of fractures, mortality, and morbidity, and is a serious threat to the quality of life of aging individuals [3].

Changes in the internal environment of the human body as it ages result in changes in bone metabolism. The amount of organic matter decreases, with a relative increase in the proportion of inorganic matter, eventually leading to osteoporosis. An increase in the fragility of the trabecular bone and a decrease in the elastic modulus cause a reduction in mechanical strength and a decrease in the load-bearing capacity of bone. Therefore, the trabecular bone is prone to microfracture, further reducing its strength and leading to the occurrence of osteoporotic fractures [4]. Osteoporosis and osteoporotic fractures create a heavy burden on society and families and has become an important public health issue worldwide.

At present, the drug treatment for primary osteoporosis mainly targets the symptoms and complications. Therapeutic mechanisms include the acceleration of bone formation, the inhibition of bone resorption, and agents that promote bone mineralization. Correcting abnormal bone turnover can improve bone mass and reduce the incidence of fractures. However, pharmacotherapy can be prohibitive in cost, take a long time to be effective, and lead to adverse reactions. Long-term use of hormone replacement therapy may increase the risk of breast cancer, stroke, and cerebral infarction [5]. Bisphosphonate therapy may lead to severe suppression of bone turnover and even osteonecrosis of the mandible [6]. Therefore, it is essential to develop improved therapies.

The fundamental mechanism of osteoporosis is an imbalance of bone resorption and formation. Excessive bone resorption leads to reduced bone mass and destruction of the bone structure [7,8]. Bergman et al. [9] indicated that osteogenesis decreased and adipogenic differentiation increased with aging in animals, and that these are the main causes of senile osteoporosis. Therefore, according to the pathogenesis of the disease, a treatment that increases the number of osteoblasts and decreases the number of adipose cells, in theory, will lead to a cure for osteoporosis.

## Bone Marrow Mesenchymal Stem Cells (BMMSCs)

The rapid development of regenerative medicine and technology for stem cell transplantation is bringing new hope to patients with osteoporosis. BMMSCs have the ability to self-renew and differentiate into a mesodermal cell line, initially derived from bone marrow stromal cells. Therefore, BMMSCs could offer a potential source of cell and gene therapy. Rodan et al. [10] demonstrated that BMMSCs have multiple differentiation potentials. Different transcription factors can regulate BMMSC differentiation into osteoblasts and adipocytes. Both of these types of cells play an important role in the maintenance of normal bone stability states.

BMMSCs do not require matching in transplantation, and do not lead to immunological rejection. Therefore, BMMSCs transplantation can be a new therapeutic method of treating osteoporosis. BMMSCs can stably express CD29, CD44, CD73, CD105, and HLA. However, some research indicated that mesenchymal stem cells do not express major histocompatibility complex (MHC) class II and do not express or have low expression levels of MHC class I. Human MHC class I is the main antigen-induced allogeneic immune rejection, resulting in mesenchymal stem cells with low immunogenicity. In addition, BMMSCs do not express T cell costimulatory molecules such as B7-1 and B7-2, and do not express or stimulate apoptosis molecules such as CD40, CD80, CD86, and FasL. Thus it lacks the second signal, which is necessary for T cell activation.

Osteoporosis is characterized by bone loss accompanied by fat tissue increase, which is an important indicator of poor hematopoietic function. In recent years, some scholars have raised the point that blood disease is one of the pathogenesises of osteoporosis. They considered that the emergence of osteoporosis is associated with chronic blood loss. Osteoporosis exists in patients with some hematological system diseases such as sickle cells anemia, chronic hemolytic anemia, and pernicious anemia [11]. There may be a relationship between osteoporosis and bone marrow hematopoietic function.

## The Mechanisms of BMMSC-Mediated Bone Regeneration

BMMSCs are separated easily, can be induced to proliferate, have multipotent differentiation potential, and retain the "homologous" feature, thus rendering them the first choice in bone tissue repair and treatment of related diseases. BMMSCs can differentiate into osteocytes, osteoblasts, adipocytes, and chondrocytes, which are widely present in bone marrow and cancellous bone and play an important role in bone metabolism [12]. Many studies have confirmed that BMMSCs can differentiate



into osteoblasts and secrete a variety of osteogenic activity factors under certain induced conditions after being isolated and cultured *in vitro*. They demonstrate good prospects in repairing bone and soft tissue injury *in vivo* [13]. Animal experiments have demonstrated that systemic or local injection of autologous BMMSCs is rarely rejected during transplantation. However, it has been reported that the capacity of BMMSCs from osteoporosis patients to differentiate into osteoblasts is lower than that in healthy individuals [14]. Rodriguez et al. [15] compared the BMMSCs from postmenopausal women with osteoporosis to those from healthy volunteers. They found that the BMMSCs from osteoporosis patients have less sensitivity to insulin-like growth factor (IGF) and a weaker ability to differentiate into osteoblasts. Zhang et al. [16] reported that there was less calcified nodule formation in the osteogenic differentiation medium, confirming the decreased osteogenic differentiation capability of BMMSCs from osteoporosis patients.

In recent years, researchers have realized the important role of bone marrow stem cells in anabolic processes in bone and have begun to fundamentally study the mechanism of abnormal BMMSCs leading to osteoporosis. Although osteoporosis is caused by a variety of mechanisms, its most basic and direct mechanism is osteoblast generation. Many other research groups have suggested other supplementary opinions. Nuttal et al. [17] considered that osteoporosis could be prevented and cured by inhibiting BMMSC differentiation into adipocytes, creating more osteoblasts. The balance of differentiation between osteoblasts and adipocytes maintains the balance of bone and adipose tissue [18]. When a large number of BMMSCs differentiate into adipocytes, the number of osteoblasts will be reduced accordingly. It is also possible that the osteoblasts and adipocytes each differentiate into the other. Even terminally differentiated cells can return to the non-stereotypical BMMSC state and then re-differentiate [19]. Therefore, the inhibition of adipogenic differentiation and promotion of osteogenic differentiation of BMMSCs have become an important method for treating osteoporosis.

## Differentiation of BMMSCs and the Treatment of Osteoporosis

### Promotion of the osteogenic differentiation of BMMSCs

A differentiation pathway of BMMSCs involves the progression from osteogenic cells into preosteoblasts and finally into osteoblasts. Once the osteoblast has formed, it can secrete several extracellular matrix proteins to control the mineralization of bone matrix. Therefore, the occurrence, proliferation, differentiation, and maturation of osteoblasts are closely related to the normal growth and development of bones, and if any one of these processes is inhibited, a bone growth

disorder results. The occurrence of osteoporosis has a direct relationship with an increase in bone resorption and a decrease in bone formation [20]. In osteoporosis treatment programs, the top priority is to improve the condition by improving osteoblast proliferation. The differentiation of BMMSCs is affected by many factors, such as hormones, cytokines, and physiotherapy. Therefore, we can treat osteoporosis by inducing BMMSCs to differentiate into osteoblasts and by promoting the number of osteoblasts.

### Therapies that promote osteogenic differentiation

#### Estrogen

The incidence of osteoporosis is significantly increased among postmenopausal women by a decrease in estrogen levels after menopause [21]. The estrogen receptor is one type of nuclear hormone receptor that has an extremely important role in maintaining the equilibrium between bone resorption and formation [22]. Gray [23] and Komm [24] indicated that the estrogen receptor is expressed in the osteoblast, and that estrogen has a regulating effect. Other studies also confirmed that estrogen has an important effect in regulating differentiation of osteoblasts [25,26]. A certain concentration of estradiol (1–10 nmol/L) can promote BMMSC proliferation, osteogenic differentiation, and chondrogenic differentiation [27]. Estrogen can also promote osteoblast proliferation, and maturation, and the expression and secretion of BMP2 [28]. Li et al. [29] indicated that estrogen enhanced the osteogenic potential and the activation of  $\beta$ -catenin in MSCs from ovariectomized rats *in vitro*. Gopalakrishnan [30] showed that estradiol has a positive effect on BMMSC-derived osteoblast proliferation and function. However, long-term hormone replacement therapy (HRT) can have adverse effects. The incidences of coronary heart disease, stroke, and breast cancer have been shown to be increased in postmenopausal women treated with estrogen.

In addition, estrogen has also been proven to inhibit adipocyte progression into fat [31]. Estrogen can act on osteoblasts or BMMSCs of the target site and inhibit the transcript levels of lipoprotein lipase, influencing the distribution of body fat. Okazaki et al. [32] indicated that estrogen not only promoted osteogenic differentiation of the BMMSCs, but also inhibited adipogenic differentiation.

### Physiotherapy that promote osteogenic differentiation

#### Pulsed electromagnetic fields (PEMFs)

Electromagnetic technology in orthopedic therapy has had more than a century of exploration. PEMFs were initially used clinically in the 1970s. In recent years, with the development



of electromagnetic technology, PEMFs have been used to promote BMMSC proliferation and differentiation [33]. Studies have shown that low-frequency PEMFs can promote osteoblast proliferation [34,35]. Simmons et al. [36] found that after stimulation with PEMFs (12 Hz, 1.1 mT), MSC proliferation was accelerated, with morphological and biological characteristics consistent with osteoblast formation. Fu et al. [37] and Sun et al. [38] indicated that PEMFs can accelerate osteogenic differentiation of cultured human BMMSCs and enhance bone repair.

Research has indicated the potential of PEMFs as a new bone treatment. It has attracted more and more interest in the field of osteoporosis [39,40]. An osteoporosis study by Takayama et al. [41] showed that PEMFs (15 Hz) can significantly increase bone mass in ovariectomized rat bone. After measuring the contents of calcium and other bone minerals, they concluded that PEMFs can significantly reduce bone loss.

Researchers have confirmed that PEMFs can promote BMMSC differentiation and proliferation by altering signaling pathways and promoting the secretion of growth factors. Through a greater understanding of PEMF mechanisms, their clinical application value will become more significant. Garland et al. [42] treated one of the knee joints of patients with chronic spinal cord injury using PEMFs for 6 months and used the other knee as the control. In the second month, the bone mineral density (BMD) of the treatment group increased by 5.1%. By the sixth month, the BMD had increased to nearly normal levels. In a study by Tabrah et al. [43] 20 patients with osteoporosis were treated with PEMFs. They found that the BMD of the radius had significantly improved, and long-term observation showed no significant changes.

Low-intensity and low-frequency PEMFs have a greater effect on osteoblast proliferation, which can accelerate cell proliferation, cell differentiation, and bone-like tissue formation. Its stimulating effect on osteoblasts occurs during the early stages of cell maturation. This may be because of the electromagnetic field acting on charged particles within the cells, especially in the small mass of the electron-ion, which then produces centripetal force, called Lorentz force. This force can regulate bone metabolism and promote BMMSC differentiation into osteoblasts, leading to increased bone formation and density. PEMFs have great potential for maintaining bone structure and increasing bone density, thereby preventing and treating osteoporosis. They show a broad range of clinical applications.

#### **Inhibition of the adipose differentiation of BMMSCs**

Osteoblasts, fat cells, and other cell lines derive from a common precursor cell – bone marrow cells. BMMSCs differentiate into osteoblasts and adipocytes with an inverse relationship.

When the BMMSCs differentiate into osteoblasts, the adipogenic differentiation is weakened; when the BMMSCs differentiate into adipocytes, the osteogenic differentiation is weakened [44,45]. Thompson et al. [46] indicated that BMMSCs can differentiate into osteoblasts and fat cells under natural conditions without any intervention, both of which maintain a dynamic equilibrium. If the balance is disrupted, a metabolic bone disease such as osteoporosis will occur. The relationship between bone formation and adipogenesis is complex in the bone marrow microenvironment. Osteoblasts and adipocytes are both derived from BMMSCs, which have pluripotent potential. There is a mutual reversal relationship and a large degree of plasticity between osteoblastic and adipogenic differentiation [47]. Clabaut et al. [48] co-cultured these 2 kinds of cells, but without direct contact. Investigation of gene expression showed that the osteoblasts tended to turn into adipocytes (lipoprotein lipase and leptin) with a decrease in the expression of osteocalcin. They concluded that the presence of adipogenic cells can contribute to osteoblast differentiation into cells with a fat cell phenotype, leading to an increase in the number of lipoblasts. This differentiation process may be triggered by the lipoblasts themselves. Therefore, for the treatment of the osteoporosis, it is essential to inhibit the proliferation of adipocytes and to promote their apoptosis. Research has shown that the number of adipocytes in the bone marrow of senile patients or postmenopausal women with osteoporosis is greater than that in healthy people, and that BMMSCs tend to differentiate into adipocytes rather than osteoblasts. For some patients with diabetes, research found that with increasing age and the administration of many hypoglycemic agents such as thiazide drugs, BMMSCs more easily differentiate into adipocytes. The number of osteoblasts decreases and that of adipocytes increases, leading to osteoporosis.

Therefore, it is very important to identify the key factors in the process of BMMSC differentiation into adipocytes. Inhibition of this process will prevent osteoporosis. This understanding will provide new targets and perspectives, generating new therapeutics. Zhao et al. [49] indicated that estrogen not only promotes BMMSC differentiation into osteoblasts but also inhibits their differentiation into adipocytes. This research provided a new theory that estrogen deficiency leads to osteoporosis. However, other groups have demonstrated that long-term estrogen replacement therapy increases the incidence of breast cancer [50,51].

The numbers of osteoblasts and adipocytes are very important for bone formation. Research has shown that glucocorticoids also have a regulatory effect on the differentiation of BMMSCs. Long-term, high-dose glucocorticoids promoted adipogenic differentiation and inhibited osteoblast differentiation, leading to osteoporosis [52]. However, in the physiological range, glucocorticoids promote BMMSC differentiation into



not only adipocytes but also osteoblasts. Low glucocorticoid concentrations favor BMMSC proliferation. However, high glucocorticoid concentrations promote lipogenesis and BMMSC proliferation, decreasing the expression of peroxisome proliferator-activated receptor gamma (PPAR $\gamma$ ) and inhibiting bone formation [53]. Kelly et al. [54] indicated that 1,25 dihydroxy vitamin D3 [1,25(OH)2D3] blocks adipogenic differentiation and inhibits lipogenesis, both induced by glucocorticoid-mediated decreased RNA expression of the late adipocyte gene markers aP2 and adipsin. Duque et al. [55] showed that 1,25(OH)2D3 also inhibits the expression of PPAR $\gamma$ 2 in senescence-accelerated mice. PPAR $\gamma$ 2 can accelerate adipogenic differentiation, which can have an important positive regulatory role during the early stage of adipogenic differentiation. Considering all of this research together, we suggest that identification of an effective method to promote osteoblast differentiation and to inhibit adipogenic differentiation will create a bright future for the treatment of osteoporosis.

### Stem Cell Transplantation and Osteoporosis

Current osteoporosis treatments target osteoclasts and osteoblasts. Although this strategy has some benefits, it does lead to long-term adverse effects. Stem cell transplantation is a new type of treatment with a unique advantage. The pathology of osteoporosis is related to a decline in the number and function of osteoblasts. We can treat osteoporosis through stem cell transplantation and promote stem cells to differentiate into osteoblasts [56]. BMMSCs are commonly used in the cell transplantation therapy field. Human BMMSCs in the body have strong potential for osteoblast differentiation and bone formation, and their transplantation into critical areas can increase bone mass, improve bone density, rebalance the internal bone tissue environment, promote osteoporotic fracture healing, and control the progress of early-stage osteoporosis. Therapeutic methods include autologous or allogeneic stem cell transplantation and gene-modified stem cell transplantation.

#### Autologous or allogeneic stem cell transplantation

Studies have reported that BMMSCs can migrate to the site of an injury, creating a favorable microenvironment for tissue repair through specific differentiation processes [57]. Animal studies have shown that autologous or allogeneic stem cell transplantation enhanced bone mass in mice with osteoporosis and that stem cells can be injected directly into the trabecular bone surface [58]. BMMSCs can participate in the repair of damaged or diseased tissue, especially in an environment in which a large number of unique cells is required. Some animal studies suggest that administering stem cells intravenously can also promote bone formation [59]. In a study by Ocarino Nde [60], 0.75 million BMMSCs isolated from healthy

rats were injected into the femurs of osteoporotic rats, and the results showed that the trabecular bone percentage in the femurs was similar to that in the femurs from healthy rats, indicating that osteoporosis may be treated by BMMSC injection. Wang et al. [61] injected BMMSCs mixed with calcium alginate gelatin into the distal femur in rabbits with osteoporosis. After 8 weeks, they showed that, compared with the injection of calcium alginate gelatin, the MSC-alginate gelatin injection group had more new bone formation, increased trabecular bone density, improved bone mechanical strength, and a strengthening effect on local bone. Okamoto et al. [62] showed that when BMMSCs were injected into the femurs of post-oophorectomy mice with osteoporosis, the bone mineral density increased significantly.

These studies represent the beginning of a new era in which autologous stem cell transplantation is becoming a potential therapeutic strategy used to treat metabolic and genetic bone diseases, including osteoporosis.

#### Cytokine improves the repair of BMMSCs

Growth and development of the tissues is a process of mutual conciliation between the cells and cytokines. Cytokines can regulate cell division, matrix synthesis, and tissue differentiation. Because of the importance of promoting tissue repair and regeneration, cytokine is considered an important element in bone tissue engineering.

The prerequisite of differentiating into osteoblasts is the proliferation of bone marrow stem cells when repairing bone injury. It is necessary to reasonably regulate the contradictions of cell proliferation and differentiation if we want to achieve an effective bone repair. Various growth factors are involved in the regulation of stem cell proliferation and differentiation, rational use of which can promote bone repair. Studies showed that bone morphogenetic protein 2 (BMP-2) is one of the strongest bone growth factors, which can induce bone formation *in vivo* and has an important role in bone tissue injury and repair [63,64]. It is also suggested that BMP-2 has a strong effect in quickly inducing BMMSCs to differentiate into osteoblasts [65]. Stromal cell-derived factor-1 (SDF-1) is the main chemokine mediating the migration of stem cells, which plays a key role in mobilization and homing of hematopoietic stem cells and progenitor cells in the outer peripheral circulation. It also can promote osteogenic differentiation of BMMSCs and stimulate angiogenesis. A series of studies showed that SDF-1 can induce BMMSCs to migrate to the damaged parts, promote osteogenic differentiation, and improve repairing efficiency [66-69]. Fibroblast growth factor 2 (FGF-2) is a bone growth factor and a potent mitogenic original, which can increase proliferative capacity of BMMSCs and produce more cell quantity, and this effect is more evident at low cell density. At



the same time, it promotes BMMSCs to differentiate into osteoblasts, promote collagen synthesis and angiogenesis, and obviously promotes osteogenesis and shortens bone tissue healing time [70,71]. In addition, FGF-2 can maintain multi-differentiation potential of MSCs [72].

### Genetically modified stem cell transplantation

Although stem cell transplantation can improve the symptoms of osteoporosis, actual treatment is limited because it is easily impeded by a variety of factors. Therefore, to improve the proliferative capacity, increase the number of transplantations, and enhance the osteogenic differentiation ability of BMMSCs, researchers hope to use genetically modified stem cells to improve therapeutic efficacy [73]. Currently, gene therapy is widely used in clinical research because of its good targeting properties. BMMSCs have become ideal carrier cells for transplantation due to their pluripotent and continuous amplification capabilities [74]. Genetically modified stem cells with altered phenotypes can change the direction of cell differentiation to inhibit adipogenic differentiation or promote osteogenic differentiation, facilitating osteogenic differentiation for the treatment of osteoporosis.

Stem cell transplantation research is opening up a new field for the treatment of bone diseases [75]. Gene therapy is also an effective and practical emerging technology. However, both have shortcomings. Cell transplantation has the limitation of prosoplasia, while gene therapy also has unavoidable disadvantages, such as immune response and expression instability. These 2 methods can complement each other, and their combination can expand the ability of stem cells to treat a greater number of diseases, but it also creates a safe, stable, and efficient expression system that does not elicit an immune response. Kumar et al. [76] transfected BMMSCs with an adenovirus carrying BMP-2 and then transplanted these cells into post-oophorectomy mice with osteoporosis. They found that osteogenic differentiation of the BMMSCs was promoted. Baltzer et al. [77] showed an impressive reduction in bone loss following the intramedullary transfer of IL-1Ra genes into ovariectomized mice. This proof of concept encouraged further development of gene therapy approaches to treat osteoporosis. In a study by Lien et al. [78], CXCR4 was transfected into mouse MSCs by adenoviral infection, yielding high expression of CXCR4. After a single intravenous infusion, they showed full recovery of bone mass and a partial restoration of bone formation in glucocorticoid-induced osteoporotic mice. Their study showed that the systemic transplantation of genetically manipulated MSCs can ameliorate osteoporosis.

In recent years, genetically modified stem cell transplantation has undergone rapid development. By combining stem cell transplantation and its associated gene expression with the

ability to promote osteogenic differentiation of BMMSCs, this method will have broad prospects for treating osteoporosis.

### Shortcomings of BMMSCs as a therapeutic agent for treating osteoporosis

It is generally accepted that BMMSCs can be cultured *in vitro*, and the chance of malignant transformation is less. However, some studies have shown that BMMSCs transplantation can promote the proliferation of tumors in some experimental models. Therefore, security and long-term adverse effects should be thoroughly investigated and verified.

There are still many issues that need to be resolved in using BMMSC transplantation to treat osteoporosis. The first issue is how to ensure the security. Tumorigenesis, immunological rejection, and the spread of pathogens are potential problems for BMMSCs transplantation [79]. Stem cells and tumor cells have many similar features, such as unlimited proliferation and strong anti-apoptotic ability [80]. Kang [81] reported that cell transplantation will contribute bone mineral metabolism disorder in the short term, and then lead to bone formation disorder and increased bone resorption. The second issue is how to maintain the potential of osteogenetic differentiation and the efficiency of bone tissue formation. However, some studies indicated that these abilities will disappear gradually over time with cell expansion *in vitro* and there will be fewer cells after implanting *in vivo* [82,83]. Multiple infusions of BMMSCs are required to improve and maintain bone density. The third issue is that the timing and environment of gene modification of stem cells must be suitable, and the level of target gene expression must be maintained *in vivo*. The fourth issue is how to choose a suitable cell scaffold material and how to promote the material compatibility and enhance the osteogenic ability of BMMSCs *in vivo*. In addition, stem cell transplantation also involves ethical issues. In a word, the microenvironment in bone marrow is complex and contains monocytes, fibroblasts, and HSCs, in addition to adipocytes and osteoblasts; these cells all play roles in aging and adipogenetic differentiation. Thus, although BMMSCs transplantation for the treatment of osteoporosis has a bright future, further research and exploration are needed.

### Conclusions

Osteoporosis is a bone disease requiring long-term treatment. With the advent of an aging society, the incidence of fractures caused by osteoporosis will significantly increase. Although the available clinically applied drugs have some effect on the treatment of osteoporosis, they are associated with adverse reactions, limiting their use. Although osteoporosis has a variety of causes, the most basic and direct mechanism is decreased



osteogenic differentiation and increased adipogenic differentiation in the bone marrow. With age, the microenvironment in the bone marrow cavity changes, increasing the number of fat cells and inhibiting bone formation. During this process, adipose tissue gradually replaces bone tissue, leading to osteoporosis. Therefore, no matter which therapeutic method is used for treating osteoporosis, it is essential to directly or indirectly promote osteogenic differentiation, increase the number of osteoblasts, inhibit fat differentiation, and improve bone metabolism, thereby treating the underlying causes of osteoporosis.

## References:

1. Raisz LG: Pathogenesis of osteoporosis: concepts, conflicts, and prospects. *J Clin Invest*, 2005; 115: 3318-25
2. Cole ZA, Dennison EM, Cooper C: Osteoporosis epidemiology update. *Curr Rheumatol Rep*, 2008; 10: 92-96
3. Xia WB, He SL, Xu L et al: Rapidly increasing rates of hip fracture in Beijing, China. *J Bone Miner Res*, 2012; 27: 125-29
4. Lane JM, Riley EH, Wiganowicz PZ: Osteoporosis: Diagnosis and treatment. *Instr Course Lect*, 1997; 46: 445-58
5. Lewiecki EM: Treatment of osteoporosis with denosumab. *Maturitas*, 2010; 66: 182-86
6. Spanou A, Lyritis GP, Chronopoulos E, Tournis S: Management of bisphosphonate-related osteonecrosis of the jaw: A literature review. *Oral Dis*, 2015; 21(8): 927-36
7. Coipeau P, Rosset P, Langonne A et al: Impaired differentiation potential of human trabecular bone mesenchymal stromal cells from elderly patients. *Cytotherapy*, 2009; 11: 584-94
8. Li C, Wei G, Gu Q et al: Proliferation and differentiation of rat osteoporosis mesenchymal stem cells (MSCs) after telomerase reverse transcriptase (TERT) transfection. *Med Sci Monit*, 2015; 21: 845-54
9. Bergman RJ, Gazit D, Kahn AJ et al: Age-related changes in osteogenic stem cells in mice. *J Bone Miner Res*, 1996; 11: 568-77
10. Rodan GA, Harada S: The missing bone. *Cell*, 1997; 89: 677-80
11. Gurevitch O, Khitrin S, Valitov A, Slavin S: Osteoporosis of hematologic etiology. *Exp Hematol*, 2007; 35: 128-36
12. Bronckers AL, Sasaguri K, Engelse MA: Transcription and immunolocalization of Runx2/Cbfa1/Peap2alphaA in developing rodent and human craniofacial tissues: Further evidence suggesting osteoclasts phagocytose osteocytes. *Microsc Res Tech*, 2003; 61: 540-48
13. Karsenty G, Ducy P, Starbuck M et al: Cbfa1 as a regulator of osteoblast differentiation and function. *Bone*, 1999; 25: 107-8
14. Rodriguez JP, Garat S, Gajardo H et al: Abnormal osteogenesis in osteoporotic patients is reflected by altered mesenchymal stem cells dynamics. *J Cell Biochem*, 1999; 75: 414-23
15. Rodriguez JP, Montecinos L, Rios S et al: Mesenchymal stem cells from osteoporotic patients produce a type I collagen-deficient extracellular matrix favoring adipogenic differentiation. *J Cell Biochem*, 2000; 79: 557-65
16. Zhang ZM, Jiang LS, Jiang SD, Dai LY: Osteogenic potential and responsiveness to leptin of mesenchymal stem cells between postmenopausal women with osteoarthritis and osteoporosis. *J Orthop Res*, 2009; 27: 1067-73
17. Nuttall ME, Gimble JM: Is there a therapeutic opportunity to either prevent or treat osteopenic disorders by inhibiting marrow adipogenesis? *Bone*, 2000; 27: 177-84
18. Beresford JM, Bennett JH, Devlin C et al: Evidence for an inverse relationship between the differentiation of adipocytic and osteogenic cells in rat marrow stromal cell cultures. *J Cell Sci*, 1992; 102 (Pt 2): 341-51
19. Gimble JM, Zvonic S, Floyd ZE et al: Playing with bone and fat. *J Cell Biochem*, 2006; 98: 251-66
20. Bouffier H, Lagneaux L, Najar M et al: The Src inhibitor dasatinib accelerates the differentiation of human bone marrow-derived mesenchymal stromal cells into osteoblasts. *BMC Cancer*, 2010; 10: 298
21. Lv H, Sun Y, Zhang Y: MiR-133 is involved in estrogen deficiency-induced osteoporosis through modulating osteogenic differentiation of mesenchymal stem cells. *Med Sci Monit*, 2015; 21: 1527-34
22. Yuan FL, Xu RS, Jiang DL et al: Leonurine hydrochloride inhibits osteoclastogenesis and prevents osteoporosis associated with estrogen deficiency by inhibiting the NF-kappaB and PI3K/Akt signaling pathways. *Bone*, 2015; 75: 128-37
23. Gray TK, Flynn TC, Gray KM, Nabell LM: 17 beta-estradiol acts directly on the clonal osteoblastic cell line UMR106. *Proc Natl Acad Sci USA*, 1987; 84: 6267-71
24. Komm BS, Terpening CM, Benz DJ et al: Estrogen binding, receptor mRNA, and biologic response in osteoblast-like osteosarcoma cells. *Science*, 1988; 241: 81-84
25. Deal C: Potential new drug targets for osteoporosis. *Nat Clin Pract Rheumatol*, 2009; 5: 20-27
26. Martin A, Xiong J, Koromila T et al: Estrogens antagonize RUNX2-mediated osteoblast-driven osteoclastogenesis through regulating RANKL membrane association. *Bone*, 2015; 75: 96-104
27. Rodriguez JP, Astudillo P, Rios S, Pino AM: Involvement of adipogenic potential of human bone marrow mesenchymal stem cells (MSCs) in osteoporosis. *Curr Stem Cell Res Ther*, 2008; 3: 208-18
28. Kim RY, Yang HJ, Song YM et al: Estrogen modulates BMP-induced sclerostin expression via the Wnt signaling pathway. *Tissue Eng Part A*, 2015; 21(13-14): 2076-88
29. Li L, Yao XL, He XL et al: Role of mechanical strain and estrogen in modulating osteogenic differentiation of mesenchymal stem cells (MSCs) from normal and ovariectomized rats. *Cell Mol Biol*, 2013; Suppl.59: OL1889-93
30. Gopalakrishnan V, Vignesh RC, Arunakaran J et al: Effects of glucose and its modulation by insulin and estradiol on BMSC differentiation into osteoblastic lineages. *Biochem Cell Biol*, 2006; 84(1): 93-101
31. Post S, Abdallah BM, Bentzon JF, Kassem M: Demonstration of the presence of independent pre-osteoblastic and pre-adipocytic cell populations in bone marrow-derived mesenchymal stem cells. *Bone*, 2008; 43: 32-39
32. Okazaki R, Inoue D, Shibata M et al: Estrogen promotes early osteoblast differentiation and inhibits adipocyte differentiation in mouse bone marrow stromal cell lines that express estrogen receptor (ER) alpha or beta. *Endocrinology*, 2002; 143: 2349-56
33. Guerkov HH, Lohmann CH, Liu Y et al: Pulsed electromagnetic fields increase growth factor release by nonunion cells. *Clin Orthop Relat Res*, 2001; (384): 265-79
34. Ongaro A, Pellati A, Bagheri L et al: Pulsed electromagnetic fields stimulate osteogenic differentiation in human bone marrow and adipose tissue derived mesenchymal stem cells. *Bioelectromagnetics*, 2014; 35: 426-36
35. Tsai MT, Li WJ, Tuan RS, Chang WH: Modulation of osteogenesis in human mesenchymal stem cells by specific pulsed electromagnetic field stimulation. *J Orthop Res*, 2009; 27: 1169-74
36. Simmons JW Jr, Mooney V, Thacker I: Pseudarthrosis after lumbar spine fusion: nonoperative salvage with pulsed electromagnetic fields. *Am J Orthop*, 2004; 33: 27-30
37. Fu YC, Lin CC, Chang JK et al: A novel single pulsed electromagnetic field stimulates osteogenesis of bone marrow mesenchymal stem cells and bone repair. *PLoS One*, 2014; 9: e91581

## Disclosure of conflict of interest

None.



38. Sun LY, Hsieh DK, Lin PC et al: Pulsed electromagnetic fields accelerate proliferation and osteogenic gene expression in human bone marrow mesenchymal stem cells during osteogenic differentiation. *Bioelectromagnetics*, 2010; 31: 209-19
39. Androjna C, Fort B, Zborowski M, Midura RJ: Pulsed electromagnetic field treatment enhances healing callus biomechanical properties in an animal model of osteoporotic fracture. *Bioelectromagnetics*, 2014; 35: 396-405
40. Shen WW, Zhao JH: Pulsed electromagnetic fields stimulation affects BMD and local factor production of rats with disuse osteoporosis. *Bioelectromagnetics*, 2010; 31: 113-19
41. Takayama K, Nomura H, Tanaka J et al: Effect of a pulsing electromagnetic field on metabolically derived osteoporosis in rats: A pilot study. *ASAIO Trans*, 1990; 36: M426-28
42. Garland DE, Adkins RH, Matsuno NN, Stewart CA: The effect of pulsed electromagnetic fields on osteoporosis at the knee in individuals with spinal cord injury. *J Spinal Cord Med*, 1999; 22: 239-45
43. Tabrah F, Hoffmeier M, Gilbert F Jr. et al: Bone density changes in osteoporosis-prone women exposed to pulsed electromagnetic fields (PEMFs). *J Bone Miner Res*, 1990; 5: 437-42
44. Yeung DK, Griffith JF, Antonio GE et al: Osteoporosis is associated with increased marrow fat content and decreased marrow fat unsaturation: A proton MR spectroscopy study. *J Magn Reson Imaging*, 2005; 22: 279-85
45. Nuttall ME, Gimble JM: Controlling the balance between osteoblastogenesis and adipogenesis and the consequent therapeutic implications. *Curr Opin Pharmacol*, 2004; 4: 290-94
46. Thompson DL, Lum KD, Nygaard SC et al: The derivation and characterization of stromal cell lines from the bone marrow of p53-/- mice: new insights into osteoblast and adipocyte differentiation. *J Bone Miner Res*, 1998; 13: 195-204
47. Verma S, Rajaratnam JH, Denton J et al: Adipocytic proportion of bone marrow is inversely related to bone formation in osteoporosis. *J Clin Pathol*, 2002; 55: 693-98
48. Clabaut A, Delplace S, Chauveau C et al: Human osteoblasts derived from mesenchymal stem cells express adipogenic markers upon coculture with bone marrow adipocytes. *Differentiation*, 2010; 80: 40-45
49. Zhao JW, Gao ZL, Mei H et al: Differentiation of human mesenchymal stem cells: the potential mechanism for estrogen-induced preferential osteoblast versus adipocyte differentiation. *Am J Med Sci*, 2011; 341: 460-68
50. Vassilopoulou-Sellin R: Breast cancer and hormonal replacement therapy. *Ann NY Acad Sci*, 2003; 997: 341-50
51. Rossouw JE, Anderson GL, Prentice RL et al: Risks and benefits of estrogen plus progestin in healthy postmenopausal women: principal results from the Women's Health Initiative randomized controlled trial. *JAMA*, 2002; 288: 321-33
52. Weinstein RS: Glucocorticoid-induced osteonecrosis. *Endocrine*, 2012; 41: 183-90
53. Zhuang H, Zhang X, Zhu C et al: Molecular mechanisms of PPAR-gamma governing MSC osteogenic and adipogenic differentiation. *Curr Stem Cell Res Ther*, 2015 [Epub ahead of print]
54. Kelly KA, Gimble JM: 1,25-Dihydroxy vitamin D3 inhibits adipocyte differentiation and gene expression in murine bone marrow stromal cell clones and primary cultures. *Endocrinology*, 1998; 139: 2622-28
55. Duque G, Macoritto M, Kremer R: 1,25(OH)2D3 inhibits bone marrow adipogenesis in senescence accelerated mice (SAM-P/6) by decreasing the expression of peroxisome proliferator-activated receptor gamma 2 (PPARGgamma2). *Exp Gerontol*, 2004; 39: 333-38
56. Zhang H, Recker R, Lee WN, Xiao GG: Proteomics in bone research. *Expert Rev Proteomics*, 2010; 7: 103-11
57. Bornes TD, Jomha NM, Mulet-Sierra A, Adesida AB: Hypoxic culture of bone marrow-derived mesenchymal stromal stem cells differentially enhances *in vitro* chondrogenesis within cell-seeded collagen and hyaluronic acid porous scaffolds. *Stem Cell Res Ther*, 2015; 6: 84
58. Ichioika N, Inaba M, Kushida T et al: Prevention of senile osteoporosis in SAMP6 mice by intrabone marrow injection of allogeneic bone marrow cells. *Stem Cells*, 2002; 20: 542-51
59. Justesen J, Stenderup K, Kassem MS: [Mesenchymal stem cells. Potential use in cell and gene therapy of bone loss caused by aging and osteoporosis]. *Ugeskr Laeger*, 2001; 163(40): 5491-95 [in Danish]
60. Ocarino Nde M, Boeloni JN, Jorgetti Y et al: Intra-bone marrow injection of mesenchymal stem cells improves the femur bone mass of osteoporotic female rats. *Connect Tiss Res*, 2010; 51: 426-33
61. Wang Z, Goh J, Das De S et al: Efficacy of bone marrow-derived stem cells in strengthening osteoporotic bone in a rabbit model. *Tissue Eng*, 2006; 12: 1753-61
62. Okamoto Y, Tateishi H, Kinoshita K et al: An experimental study of bone healing around the titanium screw implants in ovariectomized rats: Enhancement of bone healing by bone marrow stromal cells transplantation. *Implant Dent*, 2011; 20: 236-45
63. Ma XW, Cui DP, Zhao DW: Vascular endothelial growth factor/bone morphogenetic protein-2 bone marrow combined modification of the mesenchymal stem cells to repair the avascular necrosis of the femoral head. *Int J Clin Exp Med*, 2015; 8: 15528-34
64. Cheng TL, Schindeler A, Little DG: BMP-2 delivered via sucrose acetate isobutyrate (SAIB) improves bone repair in a rat open fracture model. *J Orthop Res*, 2015 [Epub ahead of print]
65. Wikesjö UM, Qahash M, Thomson RC et al: rhBMP-2 significantly enhances guided bone regeneration. *Clin Oral Implant Res*, 2004; 15: 194-204
66. Zhao T, Zhang D, Millard RW et al: Stem cell homing and angiomyogenesis in transplanted hearts are enhanced by combined intramyocardial SDF-1alpha delivery and endogenous cytokine signaling. *Am J Physiol Heart Circ Physiol*, 2009; 296(4): H976-86
67. Kimura Y, Tabata Y: Controlled release of stromal-cell-derived factor-1 from gelatin hydrogels enhances angiogenesis. *J Biomater Sci Polym Ed*, 2010; 21(1): 37-51
68. Otsuru S, Tamai K, Yamazaki T et al: Circulating bone marrow-derived osteoblast progenitor cells are recruited to the bone-forming site by the CXCR4/stromal cell-derived factor-1 pathway. *Stem Cells*, 2008; 26: 223-34
69. Kortesidis A, Zannettino A, Isenmann S et al: Stromal-derived factor-1 promotes the growth, survival, and development of human bone marrow stromal stem cells. *Blood*, 2005; 105: 3793-801
70. Boilly B, Vercoutter-Edouard AS, Hondermarck H et al: FGF signals for cell proliferation and migration through different pathways. *Cytokine Growth Factor Rev*, 2000; 11: 295-302
71. Yuan Q, Kubo T, Doi K et al: Effect of combined application of bFGF and inorganic polyphosphate on bioactivities of osteoblasts and initial bone regeneration. *Acta Biomater*, 2009; 5: 1716-24
72. Martin I, Muraglia A, Campanile G et al: Fibroblast growth factor-2 supports *ex vivo* expansion and maintenance of osteogenic precursors from human bone marrow. *Endocrinology*, 1997; 138: 4456-62
73. Parikh SN: Gene therapy: Principles and clinical applications in orthopedics. *Orthopedics*, 2004; 27: 294-303; quiz 304-5
74. Pelled G, G T, Aslan H et al: Mesenchymal stem cells for bone gene therapy and tissue engineering. *Curr Pharm Des*, 2002; 8: 1917-28
75. Centeno CJ, Busse D, Kistday J et al: Increased knee cartilage volume in degenerative joint disease using percutaneously implanted, autologous mesenchymal stem cells. *Pain Physician*, 2008; 11: 343-53
76. Kumar S, Mahendra G, Nagy TR, Ponnazhagan S: Osteogenic differentiation of recombinant adeno-associated virus 2-transduced murine mesenchymal stem cells and development of an immunocompetent mouse model for *ex vivo* osteoporosis gene therapy. *Hum Gene Ther*, 2004; 15: 1197-206
77. Baltzer AW, Whalen JD, Wooley P et al: Gene therapy for osteoporosis: evaluation in a murine ovariectomy model. *Gene Ther*, 2001; 8: 1770-76
78. Lien CY, Chih-Yuan Ho K, Lee OK et al: Restoration of bone mass and strength in glucocorticoid-treated mice by systemic transplantation of CXCR4 and cbfa-1 co-expressing mesenchymal stem cells. *J Bone Miner Res*, 2009; 24: 837-48
79. Li HC, Stoicov C, Rogers AB, Houghton J: Stem cells and cancer: evidence for bone marrow stem cells in epithelial cancers. *World J Gastroenterol*, 2006; 12: 363-71
80. Werbowetski-Ogilvie TE, Bosse M, Stewart M et al: Characterization of human embryonic stem cells with features of neoplastic progression. *Nat Biotechnol*, 2009; 27: 91-97
81. Kang MJ, Lee WY, Oh KW et al: The short-term changes of bone mineral metabolism following bone marrow transplantation. *Bone*, 2000; 26: 275-79
82. Singh SP, Chang EI, Gossain AK et al: Cyclic mechanical strain increases production of regulators of bone healing in cultured murine osteoblasts. *J Am Coll Surg*, 2007; 204: 426-34
83. Rath B, Nam J, Knobloch TJ et al: Compressive forces induce osteogenic gene expression in calvarial osteoblasts. *J Biomech*, 2008; 41: 1095-103